Stability of composite vacuum Heckmann wormholes in Brans-Dicke theory

Sergey M. Kozyrev*

Scientific center for gravity wave studies "Dulkyn", Kazan, Russia

arXiv:1401.6662v1 [gr-qc] 26 Jan 2014

Abstract

This paper discusses linearized (spherically symmetric) perturbation of static Heckmann composite thin shell wormholes in Brans-Dicke gravity. The equation of state $P = \beta^2\sigma$ at the throat is linearized around the static solution where σ energy density of the shell and P the lateral presume. We have shown that this thin shell wormholes is stable within the range $0 \leq \beta < 1$ and with all values of ω except $\omega = -2$.

PACS numbers: 04.20.-q, 04.20.Jb, 04.50.Kd

*Electronic address: Sergey@tnpko.ru

I. INTRODUCTION

After the leading work by Moris and Thorn [8], wormhole connecting two asymptotically flat spacetimes is now well known classic in relativity theories. A powerful theoretical method for constructing and analyze wormhole geometry, introduced by Visser [12, 13], is by cutting and pasting two manifolds to form a geodetsically complete one. It is well known that wormhole within the framework of general relativity always requires the presence of exotic matter [12]. On the other hand wormholes in some alternative theories of gravity have been also investigated previously and it was demonstrated that the requriment of exotic matter can be avoided.

To describe a thin shell Brans-Dicke wormhole configurations one may construct such system by two distinct strategies. Actually, the essence of the standard method [1] is the following: One takes two the same copies of spacetime manifolds with appropriate asymptotics, cuts and casts away 'useless' regions of spacetimes (containing horizons, singularities, etc.), and pastes remaining regions. As the result, one obtains a geodesically complete wormhole spacetime with given asymptotics (Schwarzschild, Reissner-Nordstrom, Brans-Dicke, etc.) and a throat being a thin shell of exotic matter violating the null energy condition. In other case, we have initially a thin shell made from *ordinary* matter, and then we look for appropriate interior and exterior Brans-Dicke solutions [1, 2] matched at the shell [11].

Here it is worth noticing that thin-shell Brans-Dicke wormholes were studied in the literature [4, 14]. Note that this approach is similar to the problem of a thin shell in general relativity (see Ref. [7]). However, the distinction is that the Birkhoff theorem is not valid in Brans-Dicke theory, and so both the interior and exterior Brans-Dicke vacua are not unique.

It is then important to know wether such wormhole are stable. Stability analysis of thin shell wormholes under small perturbations around the static solution has been done previously by several authors. In this paper, the stability of composite vacuum Brans-Dicke wormholes is analyzed under linear perturbation preserving the symmetry.

[1] The first examples of thin-shell wormholes have been given by Visser [12, 13].

II. CONSTRUCTION OF THE BRANS-DICKE COMPOSITE THIN SHELL WORMHOLES

The action of Brans-Dicke theory [5] is given by[2]

$$S = \frac{1}{2}\int dx^4\sqrt{-g}\left\{\phi R - \omega\frac{\phi^{,\mu}\phi_{,\mu}}{\phi}\right\} + S_m, \tag{1}$$

where R is the scalar curvature, ϕ is a scalar field, ω is a dimensionless coupling parameter, and S_m is an action of ordinary matter (not including the scalar field). The action (1) provides the following field equations:

$$G_{\mu\nu} = \frac{1}{\phi}T_{\mu\nu} + \frac{\omega}{\phi^2}\phi_{,\mu}\phi_{,\nu} - \frac{\omega}{2\phi^2}g_{\mu\nu}\phi_{,\alpha}\phi^{,\alpha} + \frac{1}{\phi}\phi_{;\mu;\nu} - \frac{1}{\phi}g_{\mu\nu}\phi^{;\alpha}_{;\alpha}, \tag{2a}$$

$$\phi^{;\alpha}_{;\alpha} = \frac{T}{2\omega+3}, \tag{2b}$$

where $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}$ is the Einstein tensor, and $T = T^{\mu}_{\mu}$ is the trace of the matter energy momentum tensor $T_{\mu\nu}$.

We shall consider the matching of two static and spherically symmetric spacetimes. From the mathematical point of view, the most simple and satisfactory expression for the matching conditions is, following Linchnerowicz, the assumption that there exists a system of coordinates in which the metric tensor satisfies the continuity conditions. Let ξ^i be a coordinate system on Σ where Σ is an abstract copy of any of the boundaries. Continuity conditions require a common coordinate system on Σ and this is easily done if one can set $\xi^i_+ = \xi^i_-$.

Let $(V^{\pm}, g^{\pm})$ be four-dimensional spacetimes with non-null $\Sigma^{\pm}$. The junction/shell formalism constructs a new manifold $\mathcal{M}$ by joining one of the distinct parts of V^+ to one of the distinct parts of V^- by the identification $\Sigma^+ = \Sigma^- \equiv \Sigma$. The matching conditions require the equality of the first and second fundamental forms on $\Sigma^{\pm}$. There are unique (up to orientation) unit normal vectors ${n_{\pm}}^{\alpha}$ to the boundaries. We choose them so that if ${n_+}^{\alpha}$ points towards V^+ then ${n_-}^{\alpha}$ points outside of V^- or viceversa. Clearly the sign of the normal vectors are crucial since e.g. n^-_{α} points away from the portion of V^- which will be used in forming $\mathcal{M}$. The three basis vectors tangent to Σ are

$$e^{\alpha}_i = \frac{\partial x^{\alpha}}{\partial \xi^i} \tag{3}$$

[2] Units $8\pi G = c = 1$ are used throughout the paper. Greek indices range over the coordinates of the 4-manifold and Roman indices over the coordinates of the 3-surfaces.

which give the induced metric on Σ by

$$q_{ij} = \frac{\partial x^\alpha}{\partial \xi^i}\frac{\partial x^\beta}{\partial \xi^j} g_{\alpha\beta}. \tag{4}$$

The extrinsic curvature (second fundamental form) is given by

$$\begin{aligned} K_{ij} &= \frac{\partial x^\alpha}{\partial \xi^i}\frac{\partial x^\beta}{\partial \xi^j}\nabla_\alpha n_\beta \\ &= -n_\gamma \left(\frac{\partial^2 x^\gamma}{\partial \xi^i \partial \xi^j} + \Gamma^\gamma{}_{\alpha\beta}\frac{\partial x^\alpha}{\partial \xi^i}\frac{\partial x^\beta}{\partial \xi^j} \right). \end{aligned} \tag{5}$$

Then matching conditions are simply

$$q_{ij}{}^+ = q_{ij}{}^-, \tag{6}$$

$$K_{ij}{}^+ = K_{ij}{}^-. \tag{7}$$

If both (6) and (7) are satisfied we refer to Σ as a boundary surface. If only (6) is satisfied then we refer to Σ as a thin-shell.

Writing two static spherically symmetric spacetimes V^+ and V^- with signature $(-+++)$. one can suppose that the metrics $g^+_{\alpha\beta}(x^\gamma_+)$ and $g^-_{\alpha\beta}(x^\gamma_-)$ in the coordinate systems x^γ_+ and x^γ_- are of the most general forms

$$ds^2 = -f^\pm(r,t)dt^2 + g^\pm(r,t)dr^2 + u^\pm(r,t)drdt + h^\pm(r,t)\left(d\theta^2 + \sin^2\theta d\varphi^2\right) \tag{8}$$

where $f^\pm(r,t), g^\pm(r,t), u^\pm(r,t)$ and $h^\pm(r,t)$ are of class C^2. However, in what follows, we will restrict ourselves to static spacetimes with $u^\pm(r,t) = 0$ because that is sufficient for our illustrative purpose.

Within these spacetimes define two non-null 3-surfaces Σ^+ and Σ^- with metrics $q^+_{ij}(\xi^k_+)$ and $q^-_{ij}(\xi^k_-)$ in the coordinates ξ^k_+ and ξ^k_- which decompose each of the 4-spacetimes into two distinct parts. The parametric equation for hypersurface Σ is of the form

$$F(x) = r - a = 0. \tag{9}$$

in which case the throat is located at a constant radius a. The norm to Σ can be written as

$$n_\mu = -\sqrt{g(r)f(r)}\dot{r}dt|_\Sigma + \sqrt{g(r)(1+g(r)\dot{r}^2)}dr|_\Sigma\,. \tag{10}$$

The induced metric on the Σ by the two solutions (8) is

$$q^\pm_{ij}d\xi^i_\pm d\xi^j_\pm = -f^\pm(r)d\tau^2 + h^\pm(r)\left(d\theta^2 + \sin^2\theta d\varphi^2\right) \tag{11}$$

therefore we must have on the Σ according to equality of the first fundamental form

$$f^-(a) = f^+(a), \quad h^-(a) = h^+(a) \tag{12}$$

Note that this approach not required a continuity the $g^{\pm}(a)$. Moreover, Brans-Dicke scalar field should be continuous around the shell, however, field values on the inside and the outside of the shell do not necessarily have to be same [5]. We can impose that the Σ be also characterized by equality of the second fundamental form too. This condition leads to

$$K_{11} = \frac{1}{2}\frac{f'(a)}{\sqrt{g(a)}}, \quad K_{22} = \frac{1}{2}\frac{h'(a)}{\sqrt{g(a)}} \tag{13}$$

where prime denote derivation wits respect to a. Continuity of the second fundamental form is merely equivalent to

$$h^{+'}(a)^2 g^-(a) = h^{-'}(a)^2 g^+(a), \quad f^{+'}(a)^2 g^-(a) = f^{-'}(a)^2 g^+(a) \tag{14}$$

The junction conditions in Brans-Dicke theory (generalized Darmois-Israel conditions) can be obtained by projecting on Σ the field equations (2) [10]:

$$-[K^i_j] + [K]\delta^i_j = \frac{1}{\phi}\left(S^i_j - \frac{S}{3+2\omega}\,\delta^i_j\right), \tag{15}$$

$$[\phi_{,n}] = \frac{S}{3+2\omega}, \tag{16}$$

where the notation $[Z] = Z^+|_\Sigma - Z^-|_\Sigma$ stands for the jump of a given quantity Z across the hypersurface Σ, n labels the coordinate normal to this surface and S_{ij} is the energy-momentum tensor of matter on the shell located at Σ. The quantities K and S are the traces of K^i_j and S^i_j respectively. Note that Eq. (15) is equivalent to

$$S^i_j = \phi\left(\frac{\omega+1}{\omega}[K]\delta^i_j - [K^i_j]\right). \tag{17}$$

This Lanczos equations follow from the Einstein equation for the thin shell.

III. STABILITY CONDITION (LINEARIZING METHOD).

We let the radius a to be a function of the proper time τ on the Σ and trying to find the local stability of the thin shell under small perturbation around the static solution at

$a = a_0$. The jump of the components of the extrinsic curvature associated with two sides of the hypersurface in the spacetime with the metric (8) can be found as

$$K_\tau^\tau = \frac{f(2\ddot{a}g + \dot{a}^2 g') + f'(1 + \dot{a}^2 g)\big)}{2f\sqrt{g}\sqrt{1 + \dot{a}^2 g}}, \tag{18}$$

$$K_\theta^\theta = K_\phi^\phi = \frac{h'\sqrt{g(1 + \dot{a}^2 g)}}{2gh}\,, \tag{19}$$

$$K = K_\tau^\tau + 2K_\theta^\theta. \tag{20}$$

where the overdot denotes a derivative with respect to τ.

The surface stress-energy tensor of a perfect fluid is given by

$$S_j^i = \begin{pmatrix} -\sigma & 0 & 0 \\ 0 & p & 0 \\ 0 & 0 & p \end{pmatrix}, \tag{21}$$

where σ and p are the surface energy density and the surface pressure, respectively. So $S = 2p - \sigma$. After some algebra Eq. (17) yields

$$[K_\tau^\tau] = \frac{1}{(3 + 2\omega)\phi}\left[(2 + \omega)\sigma + (2 + 2\omega)p\right] \tag{22}$$

$$\left[K_\theta^\theta\right] = -\frac{1}{(3 + 2\omega)\phi}(p + \sigma + \omega\sigma)\,. \tag{23}$$

Rearranging the Eq.(23) one can write

$$\frac{1}{2}\dot{a}^2 + V(a) = 0\,, \tag{24}$$

By solving (23) for $\dot{a}^2$, one finds

$$\dot{a}^2 = \left(\frac{2}{(3 + 2\omega)\phi}(p + \sigma + \omega\sigma)\frac{h}{[h']}\right)^2 - \frac{1}{g} \equiv -2V(a)\,. \tag{25}$$

A Taylor expansion to second order to potential $V(a)$ around the static solution yields:

$$V(a) = V(a_0) + V'(a_0)(a - a_0) + \frac{1}{2}V"(a_0)(a - a_0)^2 + O(a - a_0)^3. \tag{26}$$

Differentiating (24) with respect to τ, we have

$$V'(a) = -\ddot{a}\,. \tag{27}$$

Solving (22) for $\ddot{a}$, we obtain

$$V'(a_0) = \frac{\sqrt{1+\dot{a}^2 g}}{\sqrt{g}(3+2\omega)\phi}\left(2P(1+\omega)+\sigma(2+\omega)\right) - \frac{\dot{a}^2}{2}\left(\frac{[f']}{f}+\frac{[g']}{g}\right) - \frac{[f']}{2fg}. \tag{28}$$

It is clear that stress energy structure of the thin shell and its property are related so that an imposed equation of state will determine the stability condition. Because of its simplicity the linear equation of state $p = \beta_0^2\sigma$ is often used with the field equations for many of applications and it is frequently used in constructing thin shell. Using the parameter $\beta_0 = \frac{dp}{d\sigma}$ which is usually interpreted as a speed of sound Eq. (16) and Eq. (28) can be written as

$$V'(a_0) = -\frac{(2+\omega+2\beta^2(1+\omega))}{(-1+2\beta^2)}\frac{[\phi']}{\phi g} - \frac{\dot{a}^2}{2}\left(\frac{[f']}{f}+\frac{[g']}{g}\right) - \frac{[f']}{2fg} \tag{29}$$

Using Eq. (16) and Eq. (28) it is not difficult to see that $V(a_0) = V'(a_0) = 0$, so the potential

$$V(a) = \frac{1}{2}V"(a_0)(a-a_0)^2 + O(a-a_0)^3. \tag{30}$$

Finally, by differentiating (29) we find

$$V''(a_0) = \frac{\dot{a}^2}{2}\left(\frac{[f']^2}{f^2} - \frac{[f"]}{f} + \frac{[g']^2}{g^2} - \frac{[g"]}{g}\right) +$$
$$+\frac{(2+\omega+2\beta^2(1+\omega))}{g(2\beta^2-1)}\left(\frac{[\phi'][g']}{\phi g} + \frac{[\phi']^2}{\phi^2} - \frac{[\phi'']}{\phi}\right) + \frac{1}{2g}\left(\frac{[f']^2}{f^2} + \frac{[f'][g']}{fg} - \frac{[f'']}{f}\right) \tag{31}$$

The wormhole is stable if and only if $V''(a_0) > 0$.

IV. SPHERICALLY SYMMETRIC WORMHOLES.

The matter is concentrated at the throat, while the rest of the spacetime is the vacuum Brans-Dicke solution can be given by

$$ds^2 = -\nu_0 \left(1 - \frac{2\mu}{\rho(r)}\right)^{\frac{1}{\lambda}} dt^2 + \rho(r)'^2 \left(1 - \frac{2\mu}{\rho(r)}\right)^{-\frac{1+C}{\lambda}} dr^2 + \\ + \rho(r)^2 \left(1 - \frac{2\mu}{\rho(r)}\right)^{1-\frac{1+C}{\lambda}} d\Omega^2, \tag{32}$$

$$\phi(r) = \phi_0 \left(1 - \frac{2\mu}{\rho(r)}\right)^{\frac{C}{2\lambda}}, \tag{33}$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$ is the linear element of the unit sphere, and $\rho(r)$ the arbitrary function. Generally, this solution depends on free parameters: ϕ_0, ν_0, μ, and C. The parameter λ is not free, it obeys the following constraint condition:

$$\lambda = \sqrt{(C+1)^2 - C\left(1 - \tfrac{1}{2}\omega C\right)}. \tag{34}$$

In this solution the interpretation to a choice of function $\rho(r)$ generally correspond to a choice of state within the "interior" and "exterior" vacua. In particular, considering the function $\rho(r)$ in the form

$$\rho(r) = r, C = \frac{2}{\omega} \tag{35}$$

considerable simplification occurs, namely the interior is described by the vacuum Heckmann solution [6] :

$$f^+(r) = \alpha_0 \left(1 - \frac{B}{r}\right)^{\frac{\omega}{2+\omega}}, \tag{36a}$$

$$g^+(r) = \frac{1}{1 - \frac{B}{r}}, \tag{36b}$$

$$h^+(r) = r^2 \tag{36c}$$

$$\phi^+(r) = \phi_0 \left(1 - \frac{B}{r}\right)^{\frac{1}{2+\omega}}, \tag{36d}$$

where ϕ_0, α_0, B, and C are free (still undefined) parameters. Note that the radial coordinate r runs monotonically from B to a, where $a > B$ is a boundary of the interior region.

Assuming $C^- \equiv 0$ for exterior region of a spherical gravitating configuration one can obtain the exterior Schwarzschild solution:

$$f^-(r) = 1 - \frac{2M}{r} \tag{37a}$$

$$g^-(r) = \frac{1}{1 - \frac{2M}{r}}, \tag{37b}$$

$$h^-(r) = r^2 \tag{37c}$$

$$\phi^-(r) = 1. \tag{37d}$$

We consider a thin shell between two Heckmann manifolds with exterior parameter M (Schwarzschild mass) and interior mass parameter B. The radial coordinate r within the exterior region runs from a to infinity. We will suppose that $a > M/2$; this guarantees that the exterior region does not contain the event horizon.

So, r is the global radial coordinate monotonically running from B to a in the interior region, and from a to infinity in the exterior one. We will suppose that the all metric components and scalar field be continuous on the Σ.

$f^+(a) = f^-(a), \quad g^+(a) = g^-(a), \quad h^+(a) = h^-(a)$ and $\phi^+(a) = \phi^-(a) \equiv 1$.

Obviously, for this case Darmois-Israel conditions [9] holds. Substituting Eqs. (36) and (37) gives

$$B = M, \tag{38a}$$

$$\alpha_0 = \left(1 - \frac{2M}{a}\right)^{\frac{2}{2+\omega}}, \tag{38b}$$

$$\phi_0 = \left(1 - \frac{2M}{a}\right)^{-\frac{1}{2+\omega}}. \tag{38c}$$

At the same time, derivatives of the metric and scalar field can be discontinuous. The discontinuity of the metric is usually described in terms of a jump of the extrinsic curvature K_{ij}.

Finally, to find stable configurations, we need to compute $V''(a)$ from Eq. (31). Substituting Eq. (36) and Eq. (37) we obtain

$$V''(a_0) = 4\mu \frac{\mu\left(5 + 2\beta^2(\omega - 2) + \omega\right) - a_0\left(3 + \omega + 2\beta^2\omega\right)}{a_0^3(2\mu - a_0)\left(2\beta^2 - 1\right)\left(2 + \omega\right)} \tag{39}$$

V. STABILITY REGIONS

To find stable configurations, we need to consider all possible constraints. The wormhole is stable if and only if

$$V''(a_0) > 0. \tag{40}$$

The boundary of the region of stability is given by the surface

$$V''(a_0, \omega, \mu) = 0. \tag{41}$$

We shell now study this equation in detail. Under standard physical circumstance the shell made from a perfect fluid with the barotropic equation of state $p = \beta^2\sigma$ with $0 \leq \beta < 1$. To avoid singular behavior of the metric, the radius of the throat must satisfy

$$a_0 > 2\mu. \tag{42}$$

This conditions gives some possible cases accordingly to the value of coupling constant ω

1. Case $\omega < -2$

a. $2\mu < a_0 \leq 3\mu$, and $a_0 > 3\mu$, $\omega \leq \frac{3a_0-5\mu}{\mu-a_0}$

$$\beta > \frac{1}{\sqrt{2}}. \tag{43}$$

b. $a_0 > 3\mu$, $\frac{3a_0-5\mu}{\mu-a_0} < \omega < -2$

$$0 < \beta < \Upsilon \tag{44}$$

where $\Upsilon = \sqrt{\frac{a_0(3+\omega)-\mu(5+\omega)}{2\mu(\omega-2)-2a_0\omega}}$

2. Case $-2 < \omega < -\frac{3}{2}$

$$\Upsilon < \beta < \frac{1}{\sqrt{2}} \tag{45}$$

3. Case $\omega = -\frac{3}{2}$

$$\beta \neq \frac{1}{\sqrt{2}} \tag{46}$$

4. Case $\omega > -\frac{3}{2}$

a. $a_0 > \frac{7\mu}{3}$, $-\frac{3}{2} < \omega < \frac{2\mu}{\mu-a_0}$

$$\frac{1}{\sqrt{2}} < \beta < \Upsilon \tag{47}$$

b. $a_0 > \frac{7\mu}{3}$, $\omega \geq \frac{2\mu}{\mu-a_0}$

$$\beta > \frac{1}{\sqrt{2}} \tag{48}$$

The interior and exterior regions are matched at a thin shell made from ordinary matter with positive energy density and pressure. The resulting configuration represents a stable composite wormhole, i.e. the thin matter shell with the Schwarzschild-like exterior region and the interior region containing the wormhole throat.

VI. SUMMARY

In this paper, we have analyzed the stability of a new class of static spherically symmetric configurations composed of interior and exterior Brans-Dicke vacua divided by thin matter shell. Both vacua correspond to the same Brans-Dicke coupling parameter ω, however they are described by the solution with different sets of parameters of integration and arbitrary function $\rho(r)$. In particular, the exterior vacuum solution has $C^- \equiv 0$. In this case the solution with any ω just reduces to the Schwarzschild one being consistent with restrictions on the post-Newtonian parameters following from recent Cassini data. The interior region possesses a strong gravitational field, and so, generally, $C^+ \neq 0$.

The common geometry emerges from the junction conditions at the boundary surface. It must be combined with Darmois-Israel junction conditions. Our approach provides a clear way of showing that the interior and exterior solutions is not a unique static spherically symmetric solution.

An interesting feature of composite wormholes is that the strong-field interior region containing all exotic ghost-like matter is hidden behind the matching surface, whereas the weak-field region out of it possesses the usual Schwarzschild vacuum. Such the configuration is similar to the model of trapped-ghost wormholes [3]. Note that in both models wormholes are twice asymptotically flat. However, in the trapped-ghost wormhole model the ghost is hidden in some restricted region around the throat, whereas in the composite wormhole model the ghost-like Brans-Dicke scalar occupies the "half" of wormhole spacetime behind the matching surface [11]. Anyway, in the composite wormhole configuration a ghost is hidden in the strong-field interior region, which may in principle explain why no ghosts are observed under usual conditions.

Applying the analysis to composite wormhole geometries, considering the one of the Heckmann solutions as interior region, we deduced stability regions, and found that the latter exist well within the wide range of parameters $0 \leq \beta < 1$ and $\omega \neq -2$. We would like to remark, that the Brans-Dicke thin shell wormholes has been mentioned in [14] with a stability analysis. Actually, unlike authors claim in [14], a stability region of Heckmann composite wormholes exist with all values of ω except $\omega = -2$.

In this context the idea of linkages for vacuum regions bounding other metrics may have

broad applications.

---

[1] C. Brans and R. H. Dicke, *Mach's Principle and a Relativistic Theory of Gravitation*, Phys. Rev. **124**, 925 (1961).

[2] C. H. Brans, *Mach's Principle and a Relativistic Theory of Gravitation. II* Phys. Rev. **125**, 2194 (1962).

[3] K. A. Bronnikov and S. V. Sushkov, *Trapped ghosts: a new class of wormholes*, Class. Quant. Grav. **27**, 095022 (2010).

[4] E. F. Eiroa, M. G. Richarte, C. Simeone, *Thin-shell wormholes in BransDicke gravity*, Phys.Lett. A373 (2008) 1-4; Erratum-ibid. *Erratum to Thin-shell wormholes in BransDicke gravity*, A373 (2009) 2399-2400.

[5] Y. Fujii, N. Maeda, *The scalar-tensor theory of gravitation* (Cambridge, Cambridge University Press, 2003).

[6] O. Heckmann, P. Jordan, R.Fricke, *Zur erweiterten Gravitationstheorie I.*, Z.Astroph., 28, 113-149, (1951).

[7] A. P. Lightman, W. H. Press, R. H. Price, S. A. Teukolsky, *Problem book in relativity and gravitation* (Princeton, Princeton University Press, 1975).

[8] M.Morris, K. Thorne, U.Yurtsever, *Wormholes, Time Machines, and the Weak Energy Condition.* Phys. Rev. Let. 61, 14461449 (1988).

[9] N. Sen, Ann. Phys. (Leipzig) *Über die Grenzbedingungen des Schwerefeldes an Unstetigkeitsflächen,* **73**, 365-396 (1924); K. Lanczos, *Flächenhafte Verteilung der Materie in der Einsteinschen Gravitationstheorie*, ibid. **74**, 518540 (1924); G. Darmois, *Memorial des Sciences Mathematiques*, Fascicule XXV, Chap. V (Gauthier-Villars, Paris, 1927); W. Israel, *Singular hypersurfaces and thin shells in general relativity,* Nuovo Cimento **44B**, 1, 1-14 (1966); ibid. *Singular hypersurfaces and thin shells in general relativity,* **48B**, 2, 463 (1967).

[10] K.G. Suffern, *Singular hypersurfaces in the Brans-Dicke theory of gravity,* J. Phys. A: Math. Gen. 15, 1599 (1982); C. Barrabés and G.F. Bressange, *Singular hypersurfaces in scalar - tensor theories of gravity,* Class. Quantum Grav. 14, 805 (1997); F. Dahia and C. Romero, *Line sources in Brans-Dicke theory of gravity*, Phys. Rev. D 60, 104019 (1999).

[11] S. V. Sushkov, S. M. Kozyrev, *Composite vacuum Brans-Dicke wormholes*, Phys. Rev. D 84,

124026 .

[12] M. Visser, Phys. Rev. *Traversable wormholes: Some simple examples*, D 39, 31823184 (1989).

[13] M. Visser, *Traversable wormholes from surgically modified Schwarzschild spacetimes*, Nucl. Phys. B 328, 203-212 (1989).

[14] X. Yue, S. Gao, *Stability of Brans-Dicke thin shell wormholes*, Phys. Lett. A375, 2193-2200 (2011).